











Ministel I do 5%, Com S. ez. contains Pulsasiana Acceptable verological to 4.26 pts. Accept Tripplits. ARTERITY Services of control agreeted acids of tree bed wherein, a justic; filling with almy off he while of tight, two-replaces defects; delet; profes Jedanican broades in many, it comb take, him too tribetrountella manyers in seater, too applicate.

STRINGTED A. NET CONSERVE PL: DEC.

COPPER SULFATE BLUE VITRIOL TEC

Action ingredient: Destroyed Copper Suifele 951/4 itsel Ingredients 15/4

POISON

ANTIDOTE: Give large quantities of white of eggs or milk. Call a Physician at once.

Replace never after using. Seep container highly cleand. NET WEIGHT OUNCES PACKED BY

DGSE-Adults, 1/2 teaspoonful one or two times a day as mosted. Children, 7 to 12 years, 1/4 teaspoonful. Caution-Taking insuffers continuously may lead to a "lamping" habit. Never take a insuffur in state of monea, remitting, abdominal pains, and other sizes of appendicitie, in which case consult a physician.

DISTRIBUTED BY NET CONTENTS OZS.

ELIXIR TERPIN HYDRATE

Each II. oz. contains 0.9 gr, Codeine. Average Doses Adults, I beautoportid. Children 8-12 years, 1/2 teaspectated. To be taken every 2-4 hours when required. Do not give to younger children except

upon the advice of a physician. companied by high fever, or persists for more than a few days. (Keep bottle tightly closed)

DISTRIBUTED BY NET CONTENTS

Design: Arbith, 35 dagger Children, I deep for each year. chaid a repo. May be spread in alwesterped water on timtrop of earlies. | Keen bottle tightly risend per-

DISTRIBUTED BY HET CONTENTS FL ORS.

POISON &

DISTRIBUTED BY HET CONTINUES FL OZS

PAREGORIC

Anhydraus Marghine UE25 gm. to U.D15 gm. per 200 cc. WARRING: May be higher farming.

DOSAGE: Adults, 1 TEAspoonful one to three times I day. Child I me, ald 5 drops, I ye, ald 10 drops, 5 yes, ald 20 drops.

Not FL Ozs.

Distributed by

PEPER DRUGS

Walgreen Agency Drug Store

Phone 68

ADA, OHIO

CAMPHOR LINIMENT, N. F.

CAMPHORATED OIL FOR EXTERNAL USE ONLY.

An extend assignion for the chest, threat and tack and for sprains, strains, broken and montrolar sorrowers than to expenses, cold and fatigue CAUTICA Rasp not of reach of children.

CASTOR OIL, U. S. P.
DOSE—For adult, one tablespoonful, Children, 3 to 7
years, and transpornful, 7 to 12 years, two temporalists,
Castina—Taking incutives restlicencely may lead to a
"leastlys" habit, Never take a leastlye in cases of nauses, vemiting, abdominal pains, and other signs of

DISTRIBUTED BY NET CONTENTS PL. GES.

CAMPHOR SPIRIT, N. F.

Fig. Speciment against or severy speciment except and female, turns as a contract except and female, turns as a contract except action of the contract except actions. FOR EXTERNAL USE ONLY

Living FOR EXTENSAL USE CALLY

Section of the Process of the Section of the Section Se

Peper Drugs

PHONE 634-5881 ADA, DHID

SPIRITS TURPENTINE

IOU of Turpestinal TECHORCAL *

NOT FOR MEDICINAL USE

DISTRIBUTED BY NET CONTENTS PL. OFS.

SULFUR SUBLIMED N. F.

GLOWERS OF SULFUR

Active Ingredient: Sulfur 99.5% Inert Ingredients: not more than 0.5% For external medicinal use, technical use, or as r

NOTE-Precipitated or Washed Sulfur should be used for internal medical purposes.

Preserve in well-closed containers.

DISTRIBUTED BY NET WEIGHT.

ACETONE DANGER! EXTREMELY FLAMMABLE

Keep away from the heat and open flame. Keep oneather closed. Use with adequate ventilation, Avend prolonged or repeated contact with akin.

Technical Uses: Solvent for fats, sile, wante, rustine, rubber, pyroaylin, cellulose acetate, language, varmabas, rubber rements, nail polish remover, etc.

ELIX. LACTATED PEPSIN, N. F.

Alcohol 16 to 19%

DOSET "April, 2 temporature 3 times a day after mesh. NET FL. DES. PACKED BY













